
SUPREP AM COLONOSCOPY PREP INSTRUCTIONS

Your procedure is scheduled on: _____ Procedure Time: _____ Arrival Time: _____

Location of Procedure:	[] Central Jersey	[] RWJUH – Somerset in Somerville
	Ambulatory Surgery Center	110 Rehill Ave
	511 Courtyard Dr.	GI Center – 3 rd Flr, West Elevator
	Hillsborough, NJ 08844	Somerville, NJ 08876
	908-895-0001	908-685-2200

THE DAY BEFORE COLONOSCOPY

CLEAR LIQUIDS ONLY (see clear liquid list on next page)

** It is important to drink as much fluids as you can throughout the day to avoid dehydration and help flush the bowel.

4:00 pm

Mix 6 ounce bottle of Suprep with water in mixing container to total 16 ounces.

Drink **ALL** the liquid in the container. You must drink 32 ounces of water over the next hour.

8:00 pm

Repeat directions above for second 6 ounce bottle of Suprep. Remember to drink 32 ounces of water over the next hour.

THE DAY OF COLONOSCOPY

Continue a clear liquid diet until 4 hours prior to your procedure start time.

Take nothing by mouth within 4 hours of your procedure start time. Taking anything by mouth within 4 hours of the procedure will cause a delay or cancellation. You may take critical medications with clear liquids up until 4 hours before your procedure.

** You **MUST** have someone to drive you home

** **Cancellation Policy:** Cancellations within 48 hours incur a \$100 charge not covered by insurance.

CLEAR LIQUID LIST

Avoid all RED liquids – they can look like blood in the colon! NO ALCOHOL

- Gatorade, Powerade (Sports drinks with electrolytes are recommended to help with hydration)
- water, tea or coffee (**NO CREAM OR MILK, sugar is ok**)
- Clear Broth or Bouillon
- Jell-O, Popsicles, Italian Ice (**no fruit or cream added**)
- Apple juice, White Grape juice or White Cranberry juice, Lemonade, Ice Tea (**No juice with pulp**),
- Soda such as Sprite, 7-UP, ginger ale or any cola,
- Clear hard candy, gum.

FREQUENTLY ASKED QUESTIONS

What medications/supplements should I avoid prior to procedure?

Stop the following blood thinners, if, and only if, the Prescribing Physician approves for the number of days s/he allows: Plavix, Clopidogrel, Effient, Brillinta, Coumadin, Warfarin, Pradaxa, Xarelto and Eliquis.

What medications should I take prior to procedure?

You may take **critical** medications up until 4 hours before your start time. Do not take any vitamins/supplements the morning of your procedure. It is okay to continue Anti-inflammatory medications such as aspirin, ibuprofen, Motrin, Aleve and Tylenol.

Are there any special instructions for diabetic patients?

Hold (do not take) your oral diabetic medication the morning of the colonoscopy. Take ½ of the insulin dose prior to the procedure. Use sugar free drinks for liquids during the prep to avoid increase in blood sugar. Monitor and control your blood sugar closely during your prep to prevent low and high blood sugar. Note: Your procedure may need to be cancelled if your blood sugar is over 300mg/dl on arrival.

Are there any special instructions I need to follow in the days prior to my procedure?

Two days before the procedure it is best to avoid high fiber foods such as fresh/dried fruit, vegetables, seeds nuts, and marmalades.

Do I need to finish the entire prep if my bowel movements are clear?

Yes. It is best to finish the entire prep as directed.

What should I bring with me to my procedure?

Please bring your insurance card, photo id and current medication list.

What should I wear to my procedure?

You should wear loose comfortable clothing. Leave all jewelry and valuables at home. Contact lenses should not be worn. You should not wear any clothing with metal including zippers and underwire bras.

Are there any special instructions for breast feeding mothers or patients of child-bearing age?

Prior to procedure, breast feeding mothers must pump enough milk for 24 hours worth of feedings to have for after the procedure. They must also pump and dump for 24 hours after the procedure. Patients of child-bearing age should not empty their bladder until they have given a urine sample upon arrival.

Do I need a ride home?

You must have someone drive you home or accompany you home on public transportation. If your procedure is in the afternoon, your driver must remain at the facility during the procedure.